| Project Title | Funding | Institution | |
|--|-------------|---|--|
| Identifying gastrointestinal (GI) conditions in children with autism spectrum disorders (ASD) | \$127,500 | Harvard Medical School | |
| Neurophysiological indices of risk and outcome in autism | \$51,300 | University of Washington | |
| Neurophysiological investigation of language acquisition in infants at risk for ASD | \$28,000 | Boston University | |
| Defining high and low risk expression of emotion in infants at risk for autism | \$0 | University of Pittsburgh | |
| Observational and electrophysiological assessments of temperament in infants at risk for autism spectrum disorders | \$0 | University of Washington | |
| Early social and emotional development in toddlers at genetic risk for autism | \$373,244 | University of Pittsburgh | |
| Early identification of autism: A prospective study | \$519,453 | University of Pittsburgh | |
| Infants at risk of autism: A longitudinal study | \$599,598 | University of California, Davis | |
| Metabolic biomarkers of autism: Predictive potential and genetic susceptibility | \$351,076 | Arkansas Children's Hospital Research Institute | |
| Neurobehavioral research on infants at risk for SLI and autism | \$691,847 | Boston University Medical Campus | |
| Brain-behavior growth charts of altered social engagement in ASD infants | \$125,000 | Yale University | |
| Pupil size and circadian salivary variations in autism spectrum disorder | \$70,035 | University of Kansas | |
| Misregulation of BDNF in autism spectrum disorders | \$75,000 | Weill Cornell Medical College | |
| Model diagnostic lab for infants at risk for autism | \$599,992 | Yale University | |
| Studying the biology and behavior of autism at 1-year: The Well-Baby Check-Up approach | \$275,152 | University of California, San Diego | |
| Validation study of atypical dynamic pupillary light reflex as a biomarker for autism | \$204,525 | University of Missouri | |
| ACE Center: Integrated Biostatistical and Bioinformatic Analysis Core (IBBAC) | \$208,661 | University of California, San Diego | |
| Oxytocin biology and the social deficits of autism spectrum disorders | \$112,500 | Stanford University | |
| Physical and clinical infrastructure for research on infants-at-risk for autism at Yale | \$439,163 | Yale University | |
| Signatures of gene expression in autism spectrum disorders | \$75,000 | Children's Hospital Boston | |
| Supplement to NIH ACE Network grant: "A longitudinal MRI study of infants at risk for autism" | \$135,000 | University of North Carolina at Chapel Hill | |
| The emergence of emotion regulation in children at-risk for autism spectrum disorder | \$49,537 | University of Miami | |
| ACE Center: MRI studies of early brain development in autism | \$364,247 | University of California, San Diego | |
| ACE Network: A longitudinal MRI study of infants at risk for autism | \$3,283,233 | University of North Carolina at Chapel Hill | |
| A longitudinal 3-D MRSI study of infants at high risk for autism | \$219,046 | University of Washington | |
| ACE Center: Assessment Core | \$570,490 | Yale University | |
| RNA expression studies in autism spectrum disorders | \$250,000 | Children's Hospital Boston | |
| Are autism spectrum disorders associated with leaky-gut at an early critical period in development? | \$309,000 | University of California, San Diego | |
| Abnormal vestibulo-ocular reflexes in autism: A potential endophenotype | \$0 | University of Florida | |

| Project Title | Funding | Institution | |
|--|-----------|--|--|
| Biomarkers for autism and for gastrointestinal and sleep problems in autism | \$0 | Yale University | |
| Family/Genetic study of autism | \$70,000 | Southwestern Autism Research & Resource Center (SARRC) | |
| Development of neural pathways in infants at risk for autism spectrum disorders | \$325,029 | University of California, San Diego | |
| ACE Center: Clinical Phenotype: Recruitment and Assessment Core | \$361,993 | University of California, San Diego | |
| ACE Center: Gaze perception abnormalities in infants with ASD | \$304,365 | Yale University | |
| Identification of lipid biomarkers for autism | \$0 | Massachusetts General Hospital | |
| Multiplexed suspension arrays to investigate newborn and childhood blood samples for potential immune biomarkers of autism | \$0 | Centers for Disease Control and Prevention (CDC) | |
| Placental vascular tree as biomarker of autism/ASD risk | \$0 | Research Foundation for Mental Hygiene, Inc. | |
| Development of neural pathways in infants at risk for autism spectrum disorders (supplement) | \$244,282 | University of California, San Diego | |
| The ontogeny of social visual engagement in infants at risk for autism | \$600,325 | Yale University | |
| Developmental characteristics of MRI diffusion tensor pathway changes in autism | \$252,636 | Washington University | |
| ACE Center: Linguistic and social responses to speech in infants at risk for autism | \$304,817 | University of Washington | |
| Temporal coordination of social communicative behaviors in infant siblings of children with autism | \$28,000 | University of Pittsburgh | |
| Prospective study of infants at high risk for autism | \$292,249 | Yale University | |
| Electrophysiological, metabolic and behavioral markers of infants at risk | \$378,751 | Children's Hospital Boston | |
| Temperament, emotional expression, and emotional self-regulation in relation to later ASD diagnosis | \$0 | Bryn Mawr College | |